




**IMAGE EVALUATION
TEST TARGET (MT-3)**



**Photographic
Sciences
Corporation**

23 WEST MAIN STREET
WEBSTER, N.Y. 14580
(716) 872-4503

1.8
2.0
2.2
2.5
2.8
3.2
3.6
4.0

**CIHM/ICMH
Microfiche
Series.**

**CIHM/ICMH
Collection de
microfiches.**

01
02
03
04
05
06
07
08
09
10



Canadian Institute for Historical Microreproductions

Institut canadien de microreproductions historiques

1980

Technical and Bibliographic Notes/Notes techniques et bibliographiques

The Institute has attempted to obtain the best original copy available for filming. Features of this copy which may be bibliographically unique, which may alter any of the images in the reproduction, or which may significantly change the usual method of filming, are checked below.

L'Institut a microfilmé le meilleur exemplaire qu'il lui a été possible de se procurer. Les détails de cet exemplaire qui sont peut-être uniques du point de vue bibliographique, qui peuvent modifier une image reproduite, ou qui peuvent exiger une modification dans la méthode normale de filmage sont indiqués ci-dessous.

- Coloured covers/
Couverture de couleur
- Covers damaged/
Couverture endommagée
- Covers restored and/or laminated/
Couverture restaurée et/ou pelliculée
- Cover title missing/
Le titre de couverture manque
- Coloured maps/
Cartes géographiques en couleur
- Coloured ink (i.e. other than blue or black)/
Encre de couleur (i.e. autre que bleue ou noire)
- Coloured plates and/or illustrations/
Planches et/ou illustrations en couleur
- Bound with other material/
Relié avec d'autres documents
- Tight binding may cause shadows or distortion along interior margin/
La reliure serrée peut causer de l'ombre ou de la distorsion le long de la marge intérieure
- Blank leaves added during restoration may appear within the text. Whenever possible, these have been omitted from filming/
Il se peut que certaines pages blanches ajoutées lors d'une restauration apparaissent dans le texte, mais, lorsque cela était possible, ces pages n'ont pas été filmées.
- Additional comments:/
Commentaires supplémentaires:

- Coloured pages/
Pages de couleur
- Pages damaged/
Pages endommagées
- Pages restored and/or laminated/
Pages restaurées et/ou pelliculées
- Pages discoloured, stained or foxed/
Pages décolorées, tachetées ou piquées
- Pages detached/
Pages détachées
- Showthrough/
Transparence
- Quality of print varies/
Qualité inégale de l'impression
- Includes supplementary material/
Comprend du matériel supplémentaire
- Only edition available/
Seule édition disponible
- Pages wholly or partially obscured by errata slips, tissues, etc., have been refilmed to ensure the best possible image/
Les pages totalement ou partiellement obscurcies par un feuillet d'errata, une pelure, etc., ont été filmées à nouveau de façon à obtenir la meilleure image possible.

This item is filmed at the reduction ratio checked below/
Ce document est filmé au taux de réduction indiqué ci-dessous.

10X	12X	14X	16X	18X	20X	22X	24X	26X	28X	30X	32X
						✓					

The copy filmed here has been reproduced thanks to the generosity of:




Health Sciences Library
Queen's University

The images appearing here are the best quality possible considering the condition and legibility of the original copy and in keeping with the filming contract specifications.

Original copies in printed paper covers are filmed beginning with the front cover and ending on the last page with a printed or illustrated impression, or the back cover when appropriate. All other original copies are filmed beginning on the first page with a printed or illustrated impression, and ending on the last page with a printed or illustrated impression.

The last recorded frame on each microfiche shall contain the symbol \rightarrow (meaning "CONTINUED"), or the symbol ∇ (meaning "END"), whichever applies.

Maps, plates, charts, etc., may be filmed at different reduction ratios. Those too large to be entirely included in one exposure are filmed beginning in the upper left hand corner, left to right and top to bottom, as many frames as required. The following diagrams illustrate the method:



L'exemplaire filmé fut reproduit grâce à la générosité de:

Health Sciences Library
Queen's University

Les images suivantes ont été reproduites avec le plus grand soin, compte tenu de la condition et de la netteté de l'exemplaire filmé, et en conformité avec les conditions du contrat de filmage.

Les exemplaires originaux dont la couverture en papier est imprimée sont filmés en commençant par le premier plat et en terminant soit par la dernière page qui comporte une empreinte d'impression ou d'illustration, soit par le second plat, selon le cas. Tous les autres exemplaires originaux sont filmés en commençant par la première page qui comporte une empreinte d'impression ou d'illustration et en terminant par la dernière page qui comporte une telle empreinte.

Un des symboles suivants apparaîtra sur la dernière image de chaque microfiche, selon le cas: le symbole \rightarrow signifie "A SUIVRE", le symbole ∇ signifie "FIN".

Les cartes, planches, tableaux, etc., peuvent être filmés à des taux de réduction différents. Lorsque le document est trop grand pour être reproduit en un seul cliché, il est filmé à partir de l'angle supérieur gauche, de gauche à droite, et de haut en bas, en prenant le nombre d'images nécessaire. Les diagrammes suivants illustrent la méthode.

ails
du
odifier
une
mage

errata
to

pelure,
on à



32X

SWEDISH LECTURE

LONDON MEDICO-CHIRURGICAL SOCIETY
JUNE 16th, 1899

Etiology and Diagnosis
of Cerebro-Spinal Fever

BY

WILLIAM OSLER, M.D.

*Fellow of the Royal Society; Fellow of the Royal College of Physicians;
Professor of Medicine, Johns Hopkins University, Baltimore*

[Reprinted from the "West London Medical Journal"]



London:

JOHN BALE, SONS & DANIELSSON, LTD.
OXFORD HOUSE
93, GREAT TITCHFIELD STREET, OXFORD STREET, W.

1899



CAVENDISH LECTURE

WEST LONDON MEDICO-CHIRURGICAL SOCIETY
JUNE 16th, 1899

On the Etiology and Diagnosis
of Cerebro-Spinal Fever

BY

WILLIAM OSLER, M.D.

*Fellow of the Royal Society; Fellow of the Royal College of Physicians;
Professor of Medicine, Johns Hopkins University, Baltimore*

[Reprinted from the "West London Medical Journal"]



London:

JOHN BALE, SONS & DANIELSSON, LTD.

OXFORD HOUSE

83 89, GREAT TITCHFIELD STREET, OXFORD STREET, W.

1899

721841

no. 548

May 7/74

Schumann

\$ 45.00

Gray 7/74

Schuman

CAVENDISH LECTURE
ON
THE ETIOLOGY AND DIAGNOSIS OF
CEREBRO-SPINAL FEVER.¹

BY WILLIAM OSLER, M.D., F.R.S., F.R.C.P.
Professor of Medicine, Johns Hopkins University, Baltimore, Md.

-
-
- I. Summary of the features of cerebro-spinal fever as an epidemic.—
II. Bacteriology of cerebro-spinal fever.—III. Diagnosis.—
IV. Sporadic meningitis.—V. Note on treatment.*

IN practice we occasionally meet with a meningitis which is not tuberculous, which is not a complication of pneumonia or ulcerative endocarditis, and which is not a sequence of ear disease or of injury, and which is not a terminal infection in some chronic malady. As the meninges of both brain and cord are diffusely inflamed, we label the condition cerebro-spinal meningitis. At intervals, in certain regions, the cases multiply, and we then speak of epidemic cerebro-spinal fever. Until recently my experience, though somewhat varied, had been confined to a few instances in which the absence of the usual factors justified the diagnosis of the sporadic form of this disease. Within the past year a small outbreak in Baltimore has enabled me to study certain points in the etiology

¹ Read before the Society, June 16, 1899.

and diagnosis of this most interesting affection, and has thus determined my choice of a subject on which to address you.

I.—SUMMARY OF THE FEATURES OF CEREBRO-SPINAL FEVER AS AN EPIDEMIC.

Upon these I shall only touch very briefly.

(a) One of the most fatal of all acute diseases, it fortunately takes a very humble position among epidemics as a destroyer. No fever attacks so few individuals in a community during its periods of prevalence; so that the general mortality may be but slightly increased. On the other hand, scarcely any known fever kills so large a proportion of those attacked. The recent epidemic in Boston illustrates this very well. Of 111 cases in three hospitals 76 died, a mortality of 68·5 per cent.

(b) The outbreaks occur in certain waves or periods, of which the fourth during the present century is at present prevailing in the United States. For several years there have been local epidemics of the disease in widely separated regions, usually in villages and country districts; but in 1896, 1897 and 1898 a slight epidemic occurred in Boston, and in the latter year cases began to be recognised in Baltimore and other towns. From a recent collective investigation, made under the direction of Surgeon-General Wyman, of the U.S. Marine Hospital Service (*Public Health Reports*, vol. xiv.), we learn that cerebro-spinal fever has prevailed during the past year (as a rule in a mild form) in twenty-seven States, and in the District of Columbia. It is not possible to say that the movement of troops last summer had anything to do with the spread of the disease. The limitation for a year or more to one district or city, without spreading to adjacent towns, the localisation even (as at present in Washington and Philadelphia) to special districts of a city, the recrudescence for several seasons in succession, are well-known features illustrated by the present epidemic.

(c) Among specific diseases cerebro-spinal fever stands in some points close to pneumonia. Sporadic cases of both occur in intervals of epidemic prevalence, though in pneumonia they are much more numerous, while epidemics of pneumonia, like those of cerebro-spinal fever, are most

striking in barracks, asylums and jails. Even when not prevailing as an epidemic there may be remarkable house epidemics of cerebro-spinal fever. The seasonal relations are the same in both, and the two diseases may prevail together. The abrupt onset, the great frequency of herpes in both, the leucocytosis, "the almost identical characters of the fibrino-purulent exudate in the two diseases" (Netter), the frequent complication of pneumonia in epidemic cerebro-spinal fever and of meningitis in pneumonia are additional points of contact. The degree of contagion is about the same in both diseases, and, lastly, it has been claimed that the organism described in cerebro-spinal fever is only a variety or a degenerate form of the pneumococcus.

On the other hand, Leichtenstern, speaking against the view that the pneumococcus is the cause of epidemic cerebro-spinal fever, says:—"Pneumonia is a disease spread over the entire earth, and appears at all times, there being no land immune from it. Epidemic meningitis is very rare, and in many countries is still unknown. Croupous pneumonia attacks every age, the disposition increasing somewhat with increasing age, epidemic meningitis is a disease which affects children and young people; beyond thirty-five there is slight disposition to it. Croupous pneumonia has a typical course and a crisis; epidemic meningitis has no crisis. The complications of the two diseases are different, &c." (Quoted by Councilman, Mallory and Wright.¹)

II.—BACTERIOLOGY OF CEREBRO-SPINAL FEVER.

Until within the past few years the etiology of cerebro-spinal fever has been obscure, and a majority of observers regarded the organism found in the meningeal exudate as the pneumococcus, or a variety of it. More than twelve years ago, however, Weichselbaum described a diplococcus with special cultural peculiarities, which he regarded as the specific organism of the epidemic form of the disease. Little or no attention was paid to his communication until 1895, when his observations were confirmed by Jaeger. So little notice, indeed, had they attracted that neither Ormerod in "Allbutt's

¹ "Epidemic Cerebro-spinal Meningitis." A Report of the State Board of Health of Massachusetts. Boston, 1898.

System" (vol. i., published in 1896), nor Latimer in "Loomis and Thompson's System," published in 1897, mention Weichselbaum's organism. The work of Heubner,¹ in Germany, and more particularly the studies of Councilman, Mallory, and Wright, have fully confirmed the observations of Weichselbaum, whose organism, known variously as the meningococcus and the diplococcus intracellularis meningitidis, is now regarded as the specific cause of the disease. As the bacteriology of the subject is very fully discussed in the monograph by Councilman and his associates, and in the just issued article by Netter in vol. xvi. of the "Twentieth Century Practice," I shall confine my remarks to an account of our experience in the study of the cases which have been in my wards during the past year. To Dr. Norman B. Gwyn, who has charge of the bacteriological work in my clinical laboratory, and to Dr. N. McL. Harris, bacteriologist in the pathological department of my colleague, Prof. Welch, I am under special obligations for their careful study of the cases during life and after death.

The meningococcus in cover-slips made from the exudate has usually a diplococcus form, and lies within the polynuclear leucocytes; hence the definition *intracellularis*. Many of the cells may be stuffed with them. They may also occur free. It stains with the ordinary staining re-agents, and is decolorised by the Gram method. It grows best on Loeffler's blood serum, on which it forms "round, whitish, shining, viscid-looking colonies with smooth, sharply defined outlines, which attain a diameter of 1 to 1½ mm. in twenty-four hours" (Councilman). In cultures it is usually short-lived; but Netter states that Germano has found that it resists desiccation, and has preserved its vitality to the end even of ninety days. He states also that Neisser has shown that the organism is transportable by atmospheric currents, even the most feeble, and that it is very susceptible to aerial convection. It is found in the cerebro-spinal exudate, rarely elsewhere in the body; but it has been isolated from the blood, the pus from the joints, the pneumonic areas in the lungs and the nasal mucus.

¹ *Jahrbuch für Kinderheilkunde*, 1891, and the *Deutsche medicinische Wochenschrift*, 1897.

The organism is feebly pathogenic for animals. Rabbits and mice resist subcutaneous inoculation. Heubner, and Councilman and his associates, have produced typical meningitis by inoculating cultures of the meningococcus beneath the spinal membranes of goats.

Our clinical and pathological experience with the organism is as follows:—In twenty-one cases which I have seen, the lumbar puncture was made in sixteen. In three patients seen in consultation the diagnosis seemed so clear that lumbar puncture was not made. In cases 1 and 2, both mild, the puncture was made, in one on the seventh day, in the other on the sixth; but no organisms were found. Cases 3 and 4 were admitted late in the disease, and it was not thought necessary to perform it. Of the remaining fourteen cases, in thirteen the diplococcus intracellularis was present on cover-slips and in cultures. In the remaining case its presence was doubtful on the cover-slips, and the staphylococcus grew in culture. Of the cases which came to autopsy, five in number, the bacteriological results, as given by Dr. Harris, are as follows:—Autopsy No. 1104; meningitis, meningococcus; the pneumococcus from a pneumonic area in the lung. No. 1189, a case with extensive arthritis, in which the diplococcus intracellularis was found in the blood and joint tissues during life, as well as from the meninges. *Post-mortem*, the meningococcus was separated in pure cultures from the cord and brain, and from the lung the pneumococcus and the bacillus lactis aerogenes; from the blood in the coronary arteries the pneumococcus. The other organs were sterile. In case 10 (Autopsy No. 1247), on which laminectomy had been performed, the spinal meninges at the time of his death (about two months later) were perfectly normal. At the operation the staphylococcus pyogenes grew in the cultures. In No. 1314, a case in which laminectomy was performed, the meningococcus was isolated from the meninges, both during life and after death. The bacillus coli communis, the bacillus lactis aerogenes, and the staphylococcus pyogenes albus were also isolated; from a broncho-pneumonic area in the lung the pneumococcus and the staphylococcus pyogenes aureus, from the spleen the staphylococcus pyogenes citreus, and from a thrombus in the superior longitudinal sinus and from the pelvis of the left kidney the streptococcus septicus

liquefaciens (Sternberg). This case had a diphtheritic and hæmorrhagic cystitis. No. 1364, a very protracted case, in which death occurred in the twelfth week, and from which the meningococcus had been isolated from the spinal fluid twice, from the meninges *post-mortem* the streptococcus pyogenes and the bacillus coli communis were isolated; and the staphylococcus pyogenes aureus and the bacillus coli communis from the broncho-pneumonic areas in the lungs. In a fatal case outside the hospital the meningococcus, the pneumococcus and the bacillus coli communis grew in cultures from the cerebral meninges. There was no focus of pneumonia in the lungs.

The following memoranda by Dr. Harris are of interest:—

"Cultural methods employed. In the first cases we followed the method recommended by Councilman; namely, the use of blood serum 'slants.' Such was found to be almost useless on account of contaminating organisms, for, out of a series of eight tubes, only one gave pure cultures.

"In the other cases the exudates and fluids were plated out in alkaline 5 per cent glycerin-agar. This procedure gave entire satisfaction in that the growth of the meningococcus was quite vigorous, and that associated organisms in the exudate (Cases 1314 and 1324A) were readily separated and classified.

"Upon certain points our observations agree with those mentioned in Councilman's article.

"(a) We have never seen, as Jaeger states, a capsule formation, nor in any case unassociated with other pyogenic organisms were we able to stain the specific organism by Gram's method.

"(b) The cases coming to autopsy at the height of the disease furnished upon cover-slip and culture abundant evidence of the presence of the meningococcus, whereas cases which succumbed late in the disease yielded feeble results, or, as in case 1364, no evidence at all.

"(c) The biological features of our several cultures gave practically the same results as Councilman obtained, except that in a few cases, after carrying the organism through several generations, litmus milk was found to be slightly bleached, but with no evidence of acid formation.

"Regarding potato culture, Councilman's statement of an invisible growth on this medium should be expanded by remarking that microscopic preparations invariably show evidences of moderate increase, and that the organisms are much larger than on ordinary media.

"At no time has a blood serum culture been observed to bear any striking resemblance to a growth of pneumococcus on a similar medium.

"As in Councilman's experience, great variations in viability of the organisms was noted. But it is worthy of mention that where there is condensation water in the tube and growth present in it, the organisms as a rule will be found to be alive many days after life has ceased in the growth upon the surface of the medium.

" *Staining reactions.* Supplementary to Councilman's remark upon focal staining points when using methylene blue, it might be said that if old Loeffler's methylene blue solution is used, the coloration of that focal point is a decided violet-red.

" *Animal experiments.* Subcutaneous inoculation in rabbits and mice (white and grey house mice) was resisted in every case.

" In rabbits inoculated beneath the dura mater, or into the ventricles of the brain, death occurred in all within fourteen days to six weeks, and in no instance were we able to find evidences of gross pathological lesions, nor the presence of the meningococcus on cover-slip or culture. The animals were all greatly emaciated.

" Mice inoculated into pleural or peritoneal cavities died in from three to four days, in most cases with general septicæmia.

" *Special remarks.* Regarding Jaeger's observation of the occurrence of chains of cocci in exudate or spinal fluid, it may be said that it has not been observed, but on blood serum growths it was quite common to find chain-formation with the unstained light line of cell division running parallel to long axis of chain. However, this feature is not by any means uncommon amongst the pyogenic staphylococci.

" Attempts to isolate the specific organism from any other site than the spinal fluid, and the meninges of spinal cord and brain, have in every case been attended with failure. This is especially noteworthy in case 1189, which presented arthritis. A large quantity of the sero-purulent fluid was inoculated, incubated for three days, and the plates were found sterile, and yet cover-slips of the fluid showed considerable numbers of intracellular diplococci. The heart's blood of this case failed to give positive results at autopsy, although during life the organism had been isolated from the blood as well as from the joint lesion.

" Pneumonic or broncho-pneumonic foci have always given the pneumococcus, either alone or with the pyogenic staphylococci.

" One case, No. 1314, brings out a possible source of error in the use of Gram's stain upon the cover-slip preparations of the exudate in cases where a secondary infection may be present. The cover-slips from the meningeal exudate in this case were examined by Gram's method before cultures were ready, and cocci were found which did not decolorise, thus causing some surprise, but it was cleared of doubt by finding the staphylococcus pyogenes albus in association with the meningococcus upon the agar plate.

" Case 1325A is most interesting in that from the exudate on the cerebral meninges the meningococcus, the pneumococcus and the bacillus coli communis were isolated and positively identified. A mouse was inoculated intra-pleurally with 3cc. of a twenty-four hour old blood serum condensation-water growth, and succumbed upon the third day with a general infection.

" Another mouse received 5cc. of a strong suspension in Durham's medium of a twenty-four hour old agar-agar culture of the pneumococcus, but showed no evidence of illness. The milk culture of this organism when stained by Welch's capsule stain exhibited well-defined capsules and stained by Gram's method.

Microbic Association in Cerebro-Spinal Fever.—Councilman, Mallory and Wright state that " Mixed infections with other

organisms were not uncommon. The pneumococcus was found seven times, once in connection with Friedländer's bacillus. Terminal infections with staphylococci and streptococci were occasionally found."

Netter refers to this point in the following terms:—"In a very small number of cases observers have reported finding at the same time with the pneumococcus or the diplococcus the staphylococcus pyogenes, the streptococcus, the colon bacillus, the bacillus proteus, and the capsulated bacillus. These microbic associations are always secondary, however, and we must refrain from regarding any but the two agents above described as the primary causes of cerebro-spinal meningitis."¹

In the report on the bacteriology of the *post mortem* by Dr. Harris, the frequency and number of microbic associations will have been noticed. Three points are of interest in this connection:—In chronic cases, as 10 and 17 of our series, the diplococcus intracellularis may no longer be present; we have obtained it, however, for the first time on the 25th, 31st and 41st days of the disease. Of the two cases referred to, No. 17 presented a most typical picture of the disease, and the diplococcus intracellularis was obtained in the spinal fluid on the 31st and 35th days. At the autopsy the meninges showed a moderate amount of exudate, and there was fluid (purulent) in the posterior horns of the ventricles. The diplococcus intracellularis was not found, but the streptococcus pyogenes and the bacillus coli communis were isolated from the meninges.

Case 10 is of particular interest, and I give here a brief summary of the history:—

A sailor, John F., aged 25, was admitted October 29, 1898. The illness had begun on October 26, with swelling and pains in the right ankle, and the next day the left ankle became swollen, and then the right hip. He was able, however, to walk to the hospital. On admission there were also redness and tenderness of the left wrist. For the first week we were very doubtful as to the nature of his trouble. The temperature was between 102° and 104°, and we rather suspected typhoid fever. On the 31st he complained of pain in the back and hips, and on November 1 it was noticed that his head was in a very retracted position. On November 4 he had retention of urine. On the 6th it was noticed that he could not move his legs. Lumbar

¹ *Loc. cit.*, p. 193.

puncture was then performed, and two drachms of a thick creamy pus removed which showed cells and numbers of diplococci, some in cells, but a majority free and in groups. Dr. Cushing performed laminectomy, and drained away a large amount of purulent exudate, and irrigated the spinal meninges with normal salt solution. Cultures made from the fluid obtained by the lumbar puncture and at the operation grew the staphylococcus pyogenes aureus. From the history and symptoms the case was doubtless one of cerebro-spinal fever, subsequently infected with the staphylococcus.

More frequently the pneumococcus has been found in association with the diplococcus intracellularis. In our series it was present only once (Case 13) in the fluid obtained by lumbar puncture. On the 5th day 20 cc. of a faintly cloudy fluid were removed, and both organisms grew in culture. On the 7th day the pneumococcus was isolated from the blood. Netter found the pneumococcus associated with the diplococcus intracellularis (*post-mortem*, I take it, though it is not so stated) in ten of sixteen cases of cerebro-spinal meningitis.

A third point is the occasional association of the tubercle bacillus with the diplococcus intracellularis, but so rarely that it is not a point of much practical importance.

On the whole, then, our observations support those of Weichselbaum, Jaeger, Heubner, Councilman and others, that in epidemic cerebro-spinal fever there is an organism with special cultural peculiarities which may reasonably be regarded as the exciting cause of the disease. Netter alone, among recent observers, seems to doubt this, and says that he only found the diplococcus intracellularis in 16 out of 39 cases, and in 10 of these the pneumococcus was present at the same time. His most recent statement in the "Twentieth Century Practice" (1899) is worth quoting: "Certain observers, and more particularly Jaeger and Heubner, believe that the diplococcus intracellularis is the sole pathogenic agent in epidemic cerebro-spinal meningitis, and that the pneumococcus has no influence whatever in the production of true epidemic meningitis. Councilman is a little less emphatic, although he appears disposed to adopt this way of thinking. I do not, however, in any way accept this view. The pneumococcus can, without any doubt, cause meningitis, and in spite of Heubner's experience, the rôle of the pneumococcus has been most surely established experimentally. The

frequent occurrence of pneumonia complicating meningitis and of meningitis complicating pneumonia, and the coincidence with meningitis of many other inflammations which may be excited by the action of the pneumococcus, imply that between pneumonia and meningitis there exists a relation which cannot be denied." He seems to think that the diplococcus intracellularis may be a degenerate form of the pneumococcus, and in support of this view cites: "Its intracellular position is evidence of a phagocytosis going on, the impossibility of cultivating most of the individuals, as shown by the small number of colonies obtained, its decolorisation by Gram's method, and its usual innocuousness in animals. I would not, however, insist upon this opinion, or lay too much stress upon the arguments just mentioned, for the question is only hypothetical as yet."


It does not seem to be altogether clear from certain of Netter's observations that he has been dealing with the meningococcus. Thus in his communication (*Bull. et mem. Soc. Med. des Hop. de Paris*, xv., p. 407, 1898) the coccus which he found in the 8 cases was rounded, usually in pairs, sometimes in short chains; at other times capsules were seen. They stained by Gram's method, and in bouillon long chains developed, and mice resisted infection when inoculated subcutaneously. He concludes from these observations that the meningococcus is nothing more than a degenerate pneumococcus. Dr. Harris, in a note on this, remarks: "Such conclusions are entirely unwarranted, or if his observations are correctly stated, he evidently had the veritable pneumococcus to deal with, and by some strange mode of reasoning concludes that he has the meningococcus in his series of cases, and that it is identical with the pneumococcus, though of a degenerate form!"

Netter's position is most illogical and confusing. Throughout the article in the "Twentieth Century Practice"—in many ways one of the very best in literature—he assumes that the specific disease, cerebro-spinal fever, may be caused either by the pneumococcus or the diplococcus intracellularis. That a primary cerebro-spinal meningitis may be due to the pneumococcus is universally acknowledged; but it is in the highest degree unlikely that a remarkable specific affection like cerebro-spinal fever should be caused by two different or-

ganisms. Towards the close of the article the inconsistency of this view seems to impress him, for he says certain peculiarities "prevent us from concluding that the two diseases (*i.e.*, pneumonia and cerebro-spinal fever) are absolutely identical."

III.—DIAGNOSIS OF CEREBRO-SPINAL FEVER.

Meningitis is not always easy to recognise clinically. Mortifying *post-mortem* disclosures are even more common with it than with pericarditis. The unexpected meningitis of pneumonia, of Bright's disease, and of other chronic affections, teaches us how latent may be the process; on the other hand, who has not in typhoid fever and in pneumonia made a positive diagnosis of meningitis, and has found *post mortem* the cerebro-spinal membranes perfectly free. Sooner or later the truth of Stokes' dictum is brought home to each one of us: "There is no single nervous symptom which may not and does not occur independently of any appreciable lesion of the brain, nerves or spinal cord." The anatomical classification of lepto-meningitis is not very satisfactory, and, as I have stated, there is great lack of uniformity in the terms employed. The nomenclature of the Royal College of Physicians recognises four groups—purulent, tubercular, syphilitic, and cerebro-spinal fever. An etiological classification, possible to-day in part, is of necessity incomplete. In the annexed diagram I have made a provisional grouping of the chief forms:—



The acute primary lepto-meningitis in a large proportion of all cases follows an infection with the diplococcus intracellularis or the pneumococcus. It will be a great gain if Still's observations (to which I shall refer later) on the posterior meningitis are confirmed. I do not know where Quincke's meningitis serosa should come in an etiological grouping. Of three cases diagnosed as such in our records, in one a micococcus was isolated, but nothing further is stated as to its identification. It is highly improbable that, as Netter holds, cerebro-spinal fever either in the sporadic or epidemic forms is caused by the pneumococcus. The body of evidence accumulated within the past few years in Germany and America is strongly in favour of the specific relationship of a definite, easily recognised organism, the diplococcus intracellularis. The pneumococcus may produce a primary meningitis, and pneumonia at certain seasons and in certain regions may frequently be associated with this lesion, but both clinically and bacteriologically this form can be distinguished from the disease under consideration.

While the possibility of the occurrence must be admitted, I have not recognised a primary meningitis caused by the pyogenic organisms¹ or by the tubercle bacillus. The terminal meningitides I have placed among the secondary forms; and, as in the terminal pleurisies and peritonitides, a variety of organisms are found. Even in cerebro-spinal fever the streptococci and staphylococci, always on the watch for a *locus minoris resistentiæ*, may drive out the original foe, as in cases 10 and 17 on our list.

To the secondary forms of meningitis in the table I shall not refer, except to call attention to the importance of the pneumococcus in this group also; nor can I discuss within the limits of this lecture the clinical features of the varieties of meningitis, and must confine my remarks to a few special points in the diagnosis of cerebro-spinal fever, and to a consideration of certain sporadic forms of meningitis.

¹ The case of John Brooks (autopsy No. 1065) is of interest in this connection. He had an illness of eight weeks' duration, with obscure cerebral symptoms. On admission he had a urethral discharge. At autopsy there was found an extensive cerebral meningitis, with distension of the ventricles and slight meningitis of the dorsal cord. From the exudate and from the urethral discharge streptococci and staphylococci were isolated. The slight urethral secretion might readily have been overlooked, and the case regarded as a primary pyogenic meningitis.

I.—*Certain features in the symptomatology of cerebro-spinal fever.*
 —No other form of meningitis presents a symptomatology of the same fulness—general, cerebral, spinal, and peripheral features are present in almost every case. This results in




CHART I., CASE 7. E. R.

part from the extent of the cerebro-spinal involvement, in part from the very frequent implication of the nerves, and to some degree is a consequence of the very chronic course. I shall discuss only a few of the important symptoms.

(a) *Onset*.—In striking contrast to other forms, particularly the tuberculous, cerebro-spinal fever sets in abruptly. Without warning, while at work, or awakening the patient from a sound sleep, comes the pain in the head, &c. This peculiarity may also be met with in primary pneumococcic meningitis; but in the form secondary to pneumonia, and in that which arises in endocarditis, latency of onset is the rule.

(b) *Fever*.—There is no constant type of fever in any form of meningitis. In cerebro-spinal fever the pyrexia is very variable, and I have given here a few charts in illustration. There may be no fever at onset—as shown in Charts VIII. and IX. of the sporadic cases. Of our epidemic cases, all had fever on admission.

Irregularity in the fever may be present from the start. There is not a steady rise at first to a fastigium, but even on the third or fourth day the temperature may fall to normal. Chart I. shows this extreme irregularity in a two-hour record. The remissions were very pronounced, and the temperature rose to 108° just before death.

In other cases the course is singularly like that of typhoid fever, and for days the two-hour temperature chart may not show a variation of more than a degree. This is well illustrated in Chart II. from Case 17. This patient (as shown on the lantern slide) had the facial aspect of a case of typhoid fever. Chart III. from Case 13 shows also a very steady fever with but slight remissions to the time of death on the fifteenth day. Chart IV. from Case 15 presents a still more marked resemblance to a typhoid fever curve, particularly in the daily remissions, giving the spiked chart with which we are all so familiar in the third week of the disease. Intermissions, exacerbations, and great irregularities similar to those in tuberculosis, are very pronounced in the more chronic cases. Chart V. from Case 16 gives the maximum and minimum daily temperature from the twenty-ninth to the twenty-ninth day. It will be noticed how variable is the time of the highest point of daily fever. On thirteen days the maximum temperature was in the morning—an inverse type, to which many writers have referred, in this disease. The exacerbations are better shown in a two-hour record. In Chart VI. from Case 15 the intermittent type is shown, but the paroxysms are readily distinguished from those of tertian malarial intermittent fever by their prolonged course.

CEREBRO-SPINAL FLUID




CHART II, CASE 17. P. C.




CHART III., CASE 13. O. S.

In cases of this sort, when chills occur, the picture is very suggestive of malaria, and one can understand why some of the older writers regarded the disease as a manifestation of paludism. The chart of Case 8 is worth a passing comment. After a very steady temperature on the ninth, tenth and eleventh days the record on the twelfth day was normal. On the thirteenth day there was a paroxysm without a chill, lasting from 10 p.m. on June 19 to 4 a.m. on the 21st. On the fourteenth and fifteenth days the temperature was normal. On the evening of the latter the fever rose, and the paroxysm




CHART IV., CASE 15. J. N. M.

lasted through the sixteenth day, and at 4 p.m. the patient complained of decided chilliness. On the seventeenth day no fever. At 10 a.m. on the eighteenth day the patient had a rigor, in which the temperature rose to 104.5° ; the paroxysm lasted more than thirty hours. Then, on the twentieth, twenty-first, and twenty-second days he had recurring attacks, two with chills; the one on the twenty-second, without chill, extended into the twenty-third day. After this the patient's temperature remained normal, and he recovered rapidly. During these attacks he complained a great deal of pain in




CHART V., CASE 16. CHAS. A. K.




CHART VI., CASE 15. J. N. M.

ETIOLOGY AND DIAGNOSIS OF




CHART VII., CASE 15. C. A. K.

the head and back. The spleen was not enlarged. The blood was examined daily for the malarial parasites, but none were found. And lastly, Chart VII. from Case 15 illustrates the relation of the temperature to the lumbar puncture. After the sixth, seventh, and eighth puncture the fever dropped; after the twelfth it rose.

(c) *Skin Rashes*.—Various skin rashes are common in cerebro-spinal fever, and form an important feature in the diagnosis. Of twenty-one cases which I saw during the epidemic, in thirteen a skin eruption of some form or other was present. Herpes, of course the most common, was present in eight cases. A diffuse erythema about the chest and abdomen, and over the joints, was present in four cases. Petechiæ were noticed in eight cases, extensive in only three. In three cases a very remarkable and peculiar rash was present in the neighbourhood of the joints, particularly over the extensor surfaces of the knees and elbows, and about the ankles. There was a diffuse, livid erythema of great intensity, on which a purpuric herpes developed—a vesicular rash, the individual vesicles of which became filled with blood. As the erythema faded and the vesicles dried, they could be felt as little nodular hemispherical bodies, which persisted for a week or ten days.

(d) *Blood*.—A careful study of the blood was made in all of our cases. A leucocytosis was present in every instance. In four the first blood count was made on the third day, and the leucocytosis was 25,900, 14,500, 40,800, and 32,000 per cm. The first blood count was made on the fourth day in four cases, and the leucocytosis was 26,240, 31,800, 19,300, and 7,600 per cm. In only four cases did the leucocytes exceed 40,000 per cmm., and the maximum count was on the ninth day in a fatal case 47,000 per cm. The leucocytosis persists even in the most protracted cases. In a lad who died in the twelfth week of the disease the leucocytes were 23,000 per cm. shortly before death. As a rule there was no special reduction in the red blood corpuscles. In one case a moderate anæmia developed, and in the fifth week the red corpuscles were only 3,100,000 per cmm. In one case the diplococcus intracellularis was isolated from the blood during life.

Leucocytosis has no special value in the differential diagnosis of the various forms of meningitis. I had an impression that

it was not so constant in the tuberculous variety, but Dr. Parfitt analysed the records in eleven of our cases and found that in only three was a leucocytosis absent. The lowest count was 6,500 per cm. Of eight cases with leucocytosis there were four with 20,000 or more white corpuscles per cm. The highest count was in a child aged 8, 24,333 per cm. Altogether the leucocytosis seems not to be so high or so persistent in the tuberculous meningitis.

(c) *Arthritis*, or more often peri-arthritis, much more common in cerebro-spinal fever than in other forms, was present in two of our cases. In Case 7 the joint lesion came on with great rapidity, and by the fourth day he had a multiple suppurative arthritis, resembling an acute pyæmia. The diplococcus intracellularis was isolated from the pus in the joints. In the other (Case 10) the disease began with arthritis.¹

II.—*Kernig's Sign*.—Described by a Russian physician, and studied in Germany and France, this interesting sign has not attracted the special attention of English and American physicians, though J. B. Herrick, of Chicago, at the last meeting of the Association of American Physicians, spoke of its value. It has been present in all of our cases in which it has been looked for. It is, I think, an old observation that the subjects of protracted meningitis, particularly children, very often lie with the thighs flexed upon the abdomen, and with the legs in a state of partial contracture, so that they are with difficulty extended. To test for Kernig's sign the patient should be propped up in bed in the sitting position, then, on attempting to extend the leg on the thigh there is contraction of the flexors which prevents the full straightening of the leg. On the other hand, in the recumbent posture the leg can be fully extended. Many patients with meningitis are not in a condition to sit up, and the test can be equally well made by flexing the thigh on the abdomen, when on attempting to extend the leg, if meningitis be present, the limb cannot be fully extended, as shown in fig. 1. Friis found the sign in fifty-three of sixty cases, and Netter in forty-five of fifty. It is stated to be present in all forms of meningitis when the spinal meninges are involved. The presence of the sign is no

¹ A full report of these cases is given in the *Boston Medical and Surgical Journal*, December 29, 1898.

indication of the intensity of the spinal involvement, as it existed in a very marked degree in a recent case of pneumococcic meningitis, in which there was no positive exudate on the spinal meninges, only a turbid fluid. Netter's explanation of the phenomenon is as follows: "In consequence of the inflammation of the meninges the roots of the nerves become irritable, and the flexion of the thighs upon the pelvis when the patient is in the sitting posture elongates and consequently stretches the lumbar and sacral roots, and thus increases their irritability. The attempt to extend the knee is insufficient to provoke a reflex contraction of the flexors



FIG. 1.—Kernig's sign, showing the strong contraction of the flexors on attempting to extend the leg.

while the patient lies on his back with the thighs extended upon the pelvis, but it does so when he assumes a sitting posture."

III.—*Lumbar Puncture*.—During the past ten years no single measure of greater value in diagnosis has been introduced than Quincke's lumbar puncture. We are now able in a large number of cases to make a prompt decision as to the existence of meningitis, and are further enabled to recognise the form of the disease. I shall not detain you with details of

the technique, available now in all text-books, and recently considered at great length in an elaborate "Referat" by Neurath, in the *Centralblatt f. d. Grenzgebiete der Medizin und Chirurgie, Bd. I.* It is a simple, quite harmless procedure, and in a majority of the cases can be done without general anæsthesia, or with the aid of a local freezing mixture. A dry tap is rare in cerebro-spinal fever; the needle may be plugged with fibrin, or a nerve root may come directly against the orifice. Puncture in the third or fourth interspace may be negative, while in the second a free flow is secured. In one of our early cases in which no fluid was obtained, the autopsy showed an exudate as thick as butter, with little or no fluid. A first puncture in a case is very often negative. Though simple, the technique is, like other procedures of the kind, bettered by practice. Very often at first a few drops of blood flow, then a clear or turbid fluid, either drop by drop, or sometimes in quite a strong stream.

The fluid may be clear, turbid, purulent, or more rarely a brownish yellow or quite bloody. In a great majority of all cases when meningitis is present the fluid is turbid. In rare instances clear fluid may be obtained when meningitis exists, and in a protracted case the fluid may be turbid at one puncture and clear at the next. Several observers have noted that the fluid may become clear in the intermissions of the disease. A clear fluid may be obtained from a puncture in the second lumbar interspace, while lower down a turbid fluid may be withdrawn. I saw this possibility very well illustrated in a recent *post mortem*; the fluid in the lower dorsal and upper lumbar regions was perfectly clear, while that in the lower lumbar and the sacral regions of the canal was very turbid and contained numerous flocculent masses.

The fluid should be allowed to flow into a sterilised test-tube. When the fluid is at all turbid there is usually a slight sediment and a coagulation of fibrin. Cover-glass preparations are made either directly from the turbid fluid, or if the turbidity is slight and the cell elements few in number, after it has been centrifugalised. Cultures should be prepared at the time of making the lumbar puncture by allowing one or two cubic centimetres of the fluid to flow on to a Loeffler blood serum medium.

The amount of fluid obtained varies from a few drops to

130 cc. The samples in these flasks show the large amount which may be removed, one 126 cc., another 112 cc.

Wentworth claims that within certain limits there is some relation between the degree of turbidity of the fluid and the severity of the symptoms, and this was borne out by our more limited experience. Even in the most protracted cases the turbidity may persist. Only once was a clear fluid removed in an exacerbation. On several occasions a bloody fluid was removed in case 18. In the wards at the same time was a remarkable instance of septic meningitis in which, after turbid fluid had been withdrawn at one puncture, at a subsequent one, before death, blood flowed from the needle, and *post mortem* there was extensive meningitis and a rupture of the basilar artery with hæmorrhachis.

The number of the organisms found bears no constant relation to the intensity of the symptoms. In acute cases they are present, as a rule, in large numbers. The later the disease, the less likelihood is there to find the diplococcus intracellularis in the fluid; but we have found them in the second, third, fourth, fifth and seventh week. The following are some of the days on which the organisms were found: seventeenth day, fourteenth day, twenty-fifth day, thirty-first day, forty-first day, and nineteenth day.

Has the lumbar puncture any therapeutic value? Williams, of Boston, states that he has seen beneficial effects, and there are a few cases in the literature in which the severity of the symptoms were promptly mitigated by the removal of a variable amount of the spinal fluid. Wentworth (whose experience has been very large), speaking of these cases, says: "I have never seen any such cases, though constantly on the watch for them. A temporary relief, lasting for a few hours, has followed the operation in a few cases, but the same remissions frequently occur without any treatment." Netter says that he has seen convulsions, which had lasted for a long time without intermission, cease after the withdrawal of only about two drachms of fluid.

We have given this point our closest attention, and many times have performed the puncture directly for its supposed benefit. In case 8 the note reads: "Much better after the first puncture; brighter every way." Case 9, puncture at 11 a.m.; "Patient's condition has greatly improved; the

muttering has ceased, and the irregular movements are less marked." To case 16 I may refer more fully. It was an example of the chronic form in which the patient lingered for nearly three months. The accompanying table has been prepared by Dr. Marshall, one of my house physicians. It gives the day of the disease, the amount of fluid withdrawn, the character of the exudate, the microscopical appearances, the culture results, the leucocytosis, and the effect (?) of the puncture on the temperature. The first puncture, made on the twenty-ninth day of the illness, was negative. From the second puncture 100 cc. of a yellowish turbid fluid were withdrawn, which contained pus cells and intracellular cocci in abundance, and the cultures were positive. In all seventeen punctures were made between the twenty-ninth and the seventy-fifth day of the disease, of which fourteen were positive. A turbid pale yellow fluid was removed at each tapping. On five occasions 100 cc. or more were obtained, once 125 cc. and once 126 cc. Following the two first effective tapplings the patient appeared better, the temperature dropped, and he seemed much brighter, but he soon became worse, and the fever rose. Following the sixth, seventh, eighth, and eleventh punctures the temperature fell, 4.5°, 3.8°, 4.2°, and 5.8°. There was no change in the general condition, though he sometimes was a little brighter, and the drop in fever followed so directly that it seemed only natural to attribute the good results to the lumbar puncture. The thirteenth puncture was practically negative, yet the temperature fell 5.1°, and after the fourteenth tapping the temperature rose 2.6°. Evidently not the withdrawal of the fluid, but the peculiar character of the disease, already spoken of, was responsible for the remissions. The chart illustrates further the persistence of the leucocytosis, and the sterile characters of the fluid after the forty-fourth day. I have here for demonstration the lumbar portion of the cord and its membranes. The position of the last puncture can be seen, and the repeated operations have caused a slight hæmorrhagic pachymeningitis.

IV.—SPORADIC CEREBRO-SPINAL FEVER.

To what extent do isolated cases of cerebro-spinal fever occur in the intervals between the epidemic prevalence?

CASE XVI.—KRAITZ—LUMBAR PUNCTURE CHART.

Puncture No.	Day of Disease.	Amount.	Character.	Microscopically.	Cultures.	Leucocytosis.	Temperature.
1	29	Negative	Unchanged.
2	31	100 cc.	Yellow; turbid.	Pus cells; intracellular diplococci.	Positive.	17,400	Dropped 2°4'
3	35	125	Yellow; turbid.		Positive.	25,300	" 1°0'
4	37	90	Pale yellow; turbid.	Pus cells; no cocci.	Negative.	19,900	" 2°0'
5	40	85	Pale yellow.	Pus cells; no cocci.	"	28,000	" 1°0'
6	44	126	Pale yellow; cloudy.	Intracellular diplococci.	"	15,000	" 4°5'
7	46	100	Pale yellow; turbid.	Pus cells; no cocci.	"	22,000	" 3°8'
8	48	50	Pale yellow; turbid.	Pus cells; no cocci.	"	19,000	" 4°2'
9	54	30	Pale yellow; turbid.	Pus cells; no cocci.	"	23,600	Rose 0°8'
10	59	Negative.	Unchanged.
11	59	100 cc.	Cloudy.	Pus cells; no cocci.	"	...	Dropped 5°8'
12	62	60	Cloudy; blood-tinged.	Pus cells; no cocci.	"	18,000	" 3°4'
13	68	3	"	32,000	" 5°1'
14	70	75	Cloudy.	Pus cells; no cocci.	"	47,500	Rose 2°6'
15	73	Negative.	23,000	Unchanged.
16	75	Negative.	Unchanged.
17	75	8 cc.	Cloudy.	Pus cells; no cocci.	"	...	Unchanged.

What is the nature of the primary suppurative meningitis which is met with from time to time in all communities? Neither hospital statistics nor the ordinary death returns give trustworthy answers to these questions. The diagnosis of meningitis is often doubtful, and it is very probable that a great many cases are returned as cerebro-spinal meningitis which are the cerebral forms of typhoid fever or of pneumonia. It is surprising how much confusion there is in the diagnosis of the different forms, and the urgent need of a more accurate recognition of the varieties is well shown by the number of terms used in the tabulation of cases. Thus, not to speak of the tuberculous form, which is often spoken of as basilar meningitis, the terms most frequently used are acute meningitis, meningitis simplex, purulent meningitis, and septic meningitis. In a recently issued report on sporadic cerebro-spinal meningitis from the Philadelphia Hospital, by L. N. Boston (*Medical News*, May 20, 1899), exclusive of the tuberculous variety, the following names occur as designating the varieties of acute meningitis: leptomeningitis, basilar meningitis, pachy- and basilar meningitis, basilar and lepto-meningitis, hæmorrhagic meningitis, and purulent meningitis.

To reach uniformity in this matter is not at present possible, but we may welcome the steps which have been taken towards separating the different forms of the disease.

In the last *Census Report of the United States*, the returns for the census year 1890, a year in which cerebro-spinal fever, so far as I know, did not prevail to any extent in epidemic form, the number of deaths reported from cerebro-spinal meningitis were 3,333, a large percentage of these during the first year of life.

From the *Fifty-Ninth Annual Report of the Registrar-General*, 1896, I gather that the deaths from cerebro-spinal fever in England from 1877 to 1896 (inclusive) have only once exceeded 50 per annum. There has been a great reduction in the return since 1887, 233 cases for the ten years ending 1896, against 406 for the previous decade. In Scotland there were only six deaths from cerebro-spinal fever in 1895, and five in 1896. In Ireland there were seventy-six deaths from this cause in 1896, and the same number in 1897.

From the records of two large metropolitan hospitals I

have gleaned the following figures:—At St. Bartholomew's, from 1860, cases with the diagnosis of cerebro-spinal meningitis occur in the years 1872, 1888, 1889, 1890, 1892, 1893, 1895, 1896, 1897—twenty-one in all. In many years the word meningitis alone is used, in others the word spinal meningitis and meningitis simplex. Ormerod (*Lancet*, 1895, I.) gives an exceedingly interesting report on ten fatal cases from this Hospital: four of them occurred between March and June, 1890, at which time epidemic meningitis prevailed in the Eastern Counties. At St. Thomas's since 1873 cases with the diagnosis of cerebro-spinal meningitis occur in the years 1895 and 1896—seven cases in all. In the wards of the Royal Infirmary, Edinburgh, there were in 1891 three cases diagnosed as cerebro-spinal meningitis, in 1892 two cases, in 1893 one case.

In the United States and Canada the occurrence of sporadic cases of cerebro-spinal fever in the intervals between epidemics has long been recognised. In Montreal I performed *post mortems* on at least three cases of acute purulent cerebro-spinal meningitis not associated with pneumonia or endocarditis, and not following otitis media or injury. As illustrating the liability to error, I may mention a case in which the diagnosis of cerebro-spinal meningitis was made in malignant small-pox; in another instance, with the same diagnosis, the case was a very acute typhoid fever, fatal within a week. In Philadelphia I saw only three cases.

At the Boston City Hospital, from 1880 to 1896, there were thirty-nine cases diagnosed as cerebro-spinal meningitis with a mortality of 59 per cent. It is interesting to note that during the first five months of 1897, when the epidemic began, there were forty-two admissions, a larger number than in the previous seventeen years. That the disease lingers in a city indefinitely after an outbreak has been the common opinion of all students of the disease, and anyone who has had a large *post-mortem* experience has met with sporadic instances of extensive suppurative meningitis which he has not been able to regard as secondary to any existing condition. The Philadelphia records are interesting in this respect. As collected by Stillé from 1863 the figures illustrate very well the periods in which there were slight outbreaks; namely, 1864-65 and 1872-73. Pepper ("American Text-book of Medicine," vol. i.,

p. 163) completed the figures to 1892, and Dr. Abbott has sent me the figures up to date. They show from 1884 a progressive decline in the number of cases, which may in part have been due to more careful diagnosis. From this year, when there were 124 deaths, there was a gradual decline, and in 1891 there were only 23 deaths. From that year the figures as sent me by Dr. Abbott are as follows:—1892, 22 cases; 1893, 35 cases; 1894, 18 cases; 1895, 17 cases; 1896, 7 cases; 1897, 10 cases; 1898, 24 cases; 1899 (to and including April 31), 89 cases. Here again it is interesting to see how quickly the epidemic prevalence of the disease is manifest. During 1898 the disease was not recognised as prevalent in Philadelphia. During the present year a number of cases have occurred, and the deaths during the first four months have been 89.

In Baltimore the figures for the past six years are as follows:—1893, 59 cases; 1894, 26 cases; 1895, 27 cases; 1896, 32 cases; 1897, 22 cases; 1898, 70 cases; during this year the prevalence of the disease was recognised as occurring in a mild form through the city. During the first four months of the present year there has been an increase, and the deaths for this period have been 27.

Sporadic Cerebro-Spinal Fever at the Johns Hopkins Hospital.—In the spring of 1898 the first cases of the epidemic form were admitted to hospital, and in all eighteen cases have been under treatment. Prior to this date only four cases had been admitted which we regarded as sporadic forms of cerebro-spinal fever, three within a few days of each other in 1893, and one, a chronic case, in 1892, which has been reported.¹

CASE I.—Patrick R. (Hospital No. 7253), aged 57, admitted May 1, 1893. The patient had been drinking heavily, and when admitted was delirious, mumbling and muttering to himself, and picking at the bed-clothes, and he resisted actively any attempt to examine him. His friends stated that he had been in this condition for about a week. His temperature on admission was 97.5°, pulse 68, respirations 16. He had no skin rash and no symptoms suggestive of meningitis, and we regarded the case as one of delirium tremens. As shown in the annexed chart (Chart VIII.), he had no fever until the 5th, four days after his admission. There was then a rapid rise to 105°; the respirations increased from 28 to 40; the pulse rose to 112. Throughout the 6th the temperature remained high, falling on the morning of

¹ *Johns Hopkins Hospital Bulletin*, 1892.

the 7th to 103°. The respirations rose to 48. He was a little cyanosed, could not be roused, and he died at 8 a.m. on the 7th.

The autopsy (No. 413) showed extensive yellowish purulent exudate over the hemispheres. There was a turbid exudate at the base and along the Sylvian fissures; the ventricles contained an excess of cloudy fluid. The cord was not examined. Cover slips showed




CHART VIII. P. R.

rounded cocci in pairs, chiefly in the mono- and polynuclear cells. Cultures showed diplo-bacilli. The lungs showed slight bronchopneumonia, and the pneumococcus was isolated.

CASE 2.—Wm. F., aged 27 (Hospital No. 7282), admitted May 3, 1893. The patient was apparently very deaf, and was noisy and delirious when admitted, and he had no friends from whom any

history could be obtained. The temperature on admission was 99°, pulse 86, respirations 20. During the night he became very noisy and delirious, and the pulse was irregular. On the following morning examination of the thoracic and abdominal organs showed nothing of any special moment. The spleen was not palpable. The temperature was 99°, pulse 96, respirations 24. On the evening of the 4th




CHART IX. W. F. F.

and throughout the night the patient became very noisy and had to be transferred to the isolating ward. It was noticed on admission that he was very deaf, and this seemed to increase. On the afternoon of the 4th the temperature began to rise, and at 6 p.m. reached 102.5°. The pupils were equal; the pulse was 96, irregular. The patient resisted strongly any attempt to bend the head, and the neck was

held very rigid. There was general hyperæsthesia, and he cried out whenever he was touched. A petechial rash was noticed this afternoon for the first time over the abdomen. At 8 p.m. the temperature rose to 103° (Chart IX.). He became very much worse; the pulse became more rapid, 112, very irregular, and he died a little after 2 in the morning.

Autopsy (No. 412). A purulent greyish yellow exudate over the hemispheres, also along the pons and medulla, and the whole length of the spinal cord, which in the lumbar region reached a thickness of 8 mm. The ventricles were distended with a turbid yellowish fluid. In the posterior cornua there was a greenish exudate. Cover slips from the meninges showed numerous leucocytes and large cells containing round diplococci. Pus from the ventricles showed many cocci, single and in pairs, chiefly in cells. Some of the larger cells were packed with these cocci. Cultures were negative. Inoculations of a rabbit and a mouse were also negative. At that time we had not recognised the diplococcus intracellularis of Weichselbaum, but the description of the organism, the position in such numbers within the cells, the failure to grow in cultures and the negative results of experiments on animals, make it more than probable that we were dealing with this organism.

When these cases occurred the disease was not prevailing in Baltimore, and except in New York it was not epidemic anywhere in the country, so far as we could ascertain. Three days after the death of Case 1 a lad was admitted with a most remarkable family history of the disease. The father, a very intelligent man, gave the following details of the cases:—

Wagner family.—I. A son, a young man of about 20 years, returned home on Monday, February 7, complaining of a terrible pain in his head. He had fever with much vomiting, and his head and neck were arched. He was very delirious, became rapidly worse, and died on Saturday the 12th. II. A sister, who had helped to nurse her brother, was taken ill on Monday the 15th with similar symptoms, and died in four days. III. A second sister became affected in a few days and for two weeks was desperately ill; she then began to improve and is now well. IV. The mother, who was worn out nursing the children, was taken ill on March 17 and died in two days. She had slept in the same bed with No. V., who was admitted to the hospital.

CASE 3.—W. W., aged 15, No. V. of the Wagner family, was admitted May 12, in the eighth week of an illness which the doctor described as cerebro-spinal meningitis, and of which his mother and a brother and sister had died. The patient presented a dull heavy

appearance, and for some days did not appreciate his surroundings and appeared confused. He had delusions and hallucinations, and thought he saw his mother and sister in the room with him. He had no fever, and there was no rigidity of the head or neck. The optic nerves looked normal. He improved gradually and was discharged well June 10.

I can find no parallel to this record in the history of sporadic cerebro-spinal fever. The figures which I have given for Baltimore show that in the year 1893 a considerably larger number of cases occurred than in the four succeeding years, and the admission of these three cases in rapid succession excited our interest, but from this time until the spring of 1898 we saw nothing further of the disease. In the history of the epidemic form it is not very uncommon to find two or three cases from one house. Netter refers to Sewall's report of six children attacked in one family at the beginning of the New York epidemic in 1872, and he quotes from Thorne Thorne the statement of Morcieca that in Malta seven cases occurred in a family of nine persons. Such facts speak strongly for the contagious nature of the disease. Under ordinary circumstances, however, it is exceedingly rare to find more than one case in a family, either in the sporadic or epidemic form, and I know of no case in our recent American experience in which a physician or nurse has been attacked, or in which cases have developed among other patients in a hospital. We know nothing of the conditions under which the disease assumes this family malignancy, as it may be termed. In this respect it resembles pneumonia, not highly contagious as a rule, but which may at times, in a family or institution, present a high degree of malignancy.

Bacteriology of sporadic cerebro-spinal fever.—In a number of cases of sporadic cerebro-spinal meningitis the Weichselbaum organism has been found. The case reported by Stewart and Martin¹ is of peculiar interest, inasmuch as the epidemic form of the disease has not prevailed in Montreal for many years. It is to be noted, however, that the patient prior to her illness had been spending a few weeks in Boston, where the disease had been prevailing. An unusual point in this case was the occurrence of an acute purulent pericarditis in addition to the

¹ *Montreal Medical Journal*, March, 1898.

usual cerebro-spinal lesions. The cultures showed the diplococcus intracellularis.

By far the most suggestive contribution to this question made of late years is by Dr. George F. Still,¹ of the Great Ormond Street Hospital for Sick Children. In a study of the form of meningitis known as the simple posterior-basis of infants, he isolated from seven of eight cases a diplococcus which conforms in almost every particular with the diplococcus intracellularis. The frequency of this form of meningitis may be gathered from the fact that in ten years there were at this hospital forty-nine fatal cases, verified by autopsy. The smallest number in any one year was two, and the largest number seven. All of the cases were sporadic; none of them formed a part of any outbreak which could be considered epidemic. Clinically the disease presents certain differences from the ordinary type of cerebro-spinal fever, attacking young children, chiefly in the first year of life, and as a rule, is very much more chronic, though there are forms of cerebro-spinal fever which are quite as protracted. Skin rashes are rare, particularly herpes. The most interesting point of similarity determined by Still is the periartritic affection in certain of the cases of posterior meningitis, involving, as in epidemic form, the capsule of the joint, and the sheaths of the tendons. He determined also in one case the presence of the diplococcus in the exudate.

The problem is one which deserves the closest attention, and in every case of sporadic meningitis a careful bacteriological examination should be made for the meningococcus and the pneumococcus, as we are still ignorant of the proportion of cases due severally to these organisms. Two points remain for discussion, viz., the incidence of acute non-tuberculous meningitis in hospital work, and the clinical varieties of the pneumococcal form.

Through the kind permission of my colleague, Professor Welch, one of his assistants, Dr. W. G. MacCallum, has analysed from the pathological department of the hospital the cases of meningitis in which bacteriological examinations have been made.

The twenty-five cases may be divided into four groups:—

¹ *Journal of Pathology and Bacteriology*, vol. v., 1898.

(a) *Cerebro-spinal fever*—six cases (Nos. 412, 413, 1,104, 1,189, 1,314 and 1,362).

(b) *Pneumococcic meningitis*—eight cases (Nos. 384, 478, 619, 746, 989, 1,082, 1,151 and 1,362).

(c) *Pyogenic meningitis*, in which streptococci or staphylococci, singly or in association, were found—seven cases (Nos. 694, 1,065, 1,092, 1,247, 1,166, 1,363 and 1,364).

(d) *Miscellaneous*—four cases, two of which (Nos. 854 and 1,091) showed peculiar unidentified bacilli.

The pyogenic forms of meningitis do not concern us here; no case of primary streptococcus or staphylococcus meningitis came to autopsy. I have already referred to the chronic forms of cerebro-spinal fever in which the pyogenic cocci may alone be present at the time of death.

The meningitis with which the diplococcus lanceolatus is associated, in many respects the most important variety, deserves a separate consideration.

Pneumococcic Meningitis.

As the pneumococcus has long been recognised as the most important organism in the production of meningitis, the first question to be considered is how far sporadic cases of cerebro-spinal meningitis are due to it. Of the twenty-five cases of meningitis in the pathological department of the Johns Hopkins Hospital, in eight the pneumococcus was isolated from the exudate. Bacteriological statistics everywhere show the great importance of this organism as a factor in the causation of meningitis. Of twenty cases examined by Councilman, Mallory, and Wright there were ten cases in which the pneumococcus was found, eight of these secondary to other conditions, in two only primary. In eight cases of secondary meningitis the streptococcus was present. Netter's experience is particularly interesting. To April, 1897, he had examined bacteriologically sixty-one cases of meningitis, with the following results:—
"Pneumococcus in pure culture thirty-five times, the same associated with the streptococcus and with the staphylococcus each one time, the streptococcus alone thirteen times, the diplococcus intracellularis meningitidis three times, the staphylococcus pyogenes aureus and the bacillus of Friedländer each two times, and the coli bacillus, the influenza

bacillus, a fine bacillus, and a pyocyanic and saprophytic bacillus each one time." We may recognise three groups of cases of pneumococcic meningitis.

I. *The meningitis is a complication of lobar pneumonia.* In Montreal my attention was called to the great frequency of this association, in eight of one hundred consecutive autopsies,¹ and since these observations I have had opportunities of studying a number of cases, though I have never met with so large a proportion. By far the most important contributions to the subject have been those of Nauwerck² and Netter.³ Of the twenty-nine cases analysed by Nauwerck, in four the convexity of the brain was alone involved, in sixteen both cortex and base, and in seven the cortex, base, and spinal cord. Netter's study, the most complete that has yet appeared, contains an analysis of one hundred and twenty-four cases collected from the literature. In the recent monograph on pneumonia by Aufrecht⁴ seven cases of meningitis were found in two hundred and fifty-three autopsies, and of one thousand five hundred and one cases of pneumonia in ten the diagnosis of meningitis was made, in three of which recovery followed. Of the eight cases of pneumococcus meningitis in the department of my colleague Welch, in only four was the condition associated with lobar pneumonia.

II. *Pneumococcic meningitis from local infection.*—In this exceedingly important group the infection may come from the nose or the adjacent sinuses, from the ears, or may follow an injury to the skull, or an operation. In two of the three cases in the series analysed by Dr. MacCallum there was fracture of the base of the skull. In the third, a child, aged six, had suppuration of the nasal mucous membrane and adjacent sinuses. In this connection the importance of the pneumococcus is universally recognised, and many cases are on record of infection from an otitis media.

III. *Primary pneumococcic meningitis.*—A number of observers

¹ "The Morbid Anatomy of Pneumonia," *Canada Medical and Surgical Jour.*, May, 1885, and the Gulstonian Lectures on "Malignant Endocarditis," *British Medical Journal*, 1885, I.

² *Deutsches Archiv. für Klin. Med.*, Bd. xxix.

³ *Archives Générales de Médecine*, 1887.

⁴ Nothnagel's *Specielle Pathologie und Therapie*, Bd. xiv.

have isolated the pneumococcus in primary suppurative meningitis. The records cannot be regarded, however, as entirely trustworthy, since it is only within the past few years that bacteriologists have learned to recognise the differences between the pneumococcus and the diplococcus intracellularis. Subsequent study must determine in what proportion of cases of sporadic meningitis the pneumococcus exists, and in how many the diplococcus intracellularis.

The primary pneumococcic meningitis may develop in a person in good health, but more frequently there is a general pneumococcus infection in a debilitated individual, or in one with chronic disease. The meningitis may be present alone, or as is so common, in association with endocarditis. Of the cases examined in the *post-mortem* room of the Johns Hopkins hospital, one (No. 989), a man aged 44, had a general infection with the pneumococcus and meningitis. Another case (No. 1,206), with a general infection, acute endocarditis, and meningitis, is not included in this series, as a partial autopsy only was allowed, and but a small portion of the lumbar cord was examined.

Clinical features of pneumococcic meningitis.—I shall now refer to certain clinical features of these three groups of cases:—

(a) *Pneumonia with complicating meningitis.*—Is the case one of cerebro-spinal fever with pneumonia or of inflammation of the lungs with an added meningitis? This question does not often arise now at the bed-side, as it is most exceptional for the meningitis of pneumonia to present the symptoms of cerebro-spinal fever, and in a dubious case occurring during an epidemic the lumbar puncture (as in a case to be referred to in a few minutes) may be relied upon to clear up any doubts. The most valuable clinical record of meningitis in pneumonia is to be found in Nauwerck's paper. The histories, seventeen in number, are very full and complete, and in every case accompanied with a *post-mortem* report. He has added a series of twelve cases from the literature, making twenty-nine in all. All of the cases were above the twentieth year of age—a striking contrast to cerebro-spinal fever, in which a large proportion of all the cases are in the young. A second point is the latency of meningitis in pneumonia, which is much more often recognised in the dead house than in the wards. Netter states that fully one-half of the cases are of

this latent type. Of the cases I saw in Montreal I remember but one in which the diagnosis was made during life. Huguenin, referring to this obscurity, says: "We know no symptom which is constant in all cases or which may not be present in an identical manner in other conditions." Headache, early delirium, deepening into unconsciousness, are present in all cases. This is a consequence of the more common involvement of the cortex of the hemispheres. As Leichtenstern remarks, the mind may remain clear throughout the course of a case of cerebro-spinal fever. Spinal symptoms are rare; in only seven of the cases analysed by Nauwerck was there rigidity of the neck muscles. Strabismus was present in one-fifth of the cases, ptosis only once. The importance of lumbar puncture cannot be too strongly emphasised. A recent case, which occurred while cases of cerebro-spinal fever were in the hospital, illustrates this so well that I give an abstract of it:—

Charles E., aged 52 (Hospital No. 26417), was admitted May 6, 1899, with pneumonia of the right upper lobe of about eight days' duration. He was in a semi-comatose condition, with a temperature of only 99° , respirations 32, pulse 108. There was a pneumonia of the apex of the right lung. The day after admission it was noted that his arms and legs were very stiff; Kernig's sign was well marked, and there was some rigidity of the neck. The patient, however, responded to questions. Lumbar puncture was made in the third interspace, and 5 cc. of a perfectly clear limpid fluid obtained. Lumbar puncture was made again on May 10, and 30 cc. of a perfectly clear fluid obtained, negative on cover-slips and culture. For the next three days he had a good deal of tremor. Kernig's sign was present, and there was still some rigidity of the arms and of the neck. The leucocytes ranged from 16,000 to 24,000 per cmm. On the 14th lumbar puncture was performed for a third time, and a slightly turbid fluid obtained, from which on cover-slips the pneumococcus was obtained. The patient died on the evening of the 14th. The autopsy showed a pneumonia of the right apex, a very slight purulent exudate at the base, some turbid fluid in the ventricles, and a small amount of turbid cerebro-spinal fluid, but no positive exudate on the spinal meninges. There was an ulcerative endocarditis of the aortic valve. The pneumococcus was isolated from the meninges, the heart valves, and from the lung. A very interesting feature in this case was the normal temperature on the 11th, 12th and 13th days of the disease, and the high temperature on the 14th and 15th days, ranging to 105° and 106° .

And lastly, a most important difference between the meningitis complicating pneumonia and cerebro-spinal fever is the almost universally fatal course of the former. Nauwerck

describes the complication as invariably fatal, and he was not able to convince himself that in any one of the reported cases of recovery in the literature meningitis was actually present. Netter¹ speaks somewhat more hopefully, and states that recovery may occur. He gives two cases from the literature, both in children. In one, a child of five, in which the crisis occurred on the sixth day; the signs of meningitis were the presence during the first few days of headache, unequal pupils and ptosis. Aufrecht, in his recent monograph, *loc. cit.*, also speaks of the possibility of recovery. Of ten cases of meningitis in 1501 of pneumonia, three recovered. Netter, in his article in "the Twentieth Century Practice," commenting upon the statement by Wentworth that pneumococcal meningitis is always fatal, states that "as early as 1887 I demonstrated, however, that its curability is one of the most evident characteristics of the pneumococcal meningitis." So far as I can gather there is nothing in his essay to justify such a statement, since, as I have mentioned, he gives but two instances of recovery.

Personally I have never recognised recovery in pneumonia complicated with meningitis, and the literature of the subject bears out strongly the view that one of the most striking differences between the meningitis of pneumonia and cerebro-spinal fever is the almost invariably fatal termination of the former.

(b) *Pneumococcal meningitis from local infection.* — Two of the three cases in this group were surgical; one (No. 746), followed removal of the bones of nose and orbit; the other (1151) a fracture of the base of the skull, involving the cribriform plate of the ethmoid. In both the pneumococcus alone was isolated from the exudate in the cerebral meningitis. In two of Councilman's eight cases the infection followed fracture, in one otitis media. Cases of this group do not so often appear in the medical wards, but last year we were particularly interested in a case in which the infection started from the nose and accessory sinuses; though the possibility has to be considered of extension from the meninges. An abstract is worth quoting, and the picture (fig. 2) shows a remarkable condition of cervical opisthotonos:

¹ *Archives Generales de Medicine*, 1887, I., p. 273.

Anna C., aged 6 (black), admitted March 31, 1898. Her previous history was negative. Five days before entrance, directly after supper, she vomited and complained of headache; later in the evening she became delirious. Two days after the head was held stiffly, and she had much pain in the back of the neck. She had fever, rapid pulse, loss of appetite, and constipation. On admission she was conscious, but very irritable and difficult to examine. The head was arched back and could not be brought forward. The pupils were equal and reacted to light. There was a marked slight internal strabismus. There was nothing of moment found in the examination of the chest and abdomen. She was very tender in the sub-occipital region, but there was no sign of disease of the cervical spine. Lumbar puncture on the 1st was negative. For the first week the condition remained about the same; the child could be roused, and answered



FIG. 2.—A. C. Pneumococcic meningitis—cervical opisthotonos.

questions, the strabismus persisted; there were no changes in the optic nerves. The temperature ranged from 100° to 104° . The arching of the head and neck continued; the eyes looked directly upwards (fig. 2). On the 17th a rash appeared, a diffuse erythema over the face, neck and chest, with papules. The rash faded in three days. The temperature ranged from 102° to 104° . On April 19 a profuse purulent discharge appeared from the nose, which showed the pneumococcus on cover-slips, but in cultures only the staphylococcus aureus grew. She could still be roused; the attitude remained as shown in the photograph. No optic neuritis. Throughout the 18th the temperature dropped from 105° to 97° on the morning of the 19th, and continued about 97° until her death on the morning of the 21st. One of the most remarkable features was the retention of consciousness to within a week of her death.

The autopsy (No. 1082) showed exudate over the sulci of the hemispheres. The base, from just behind the optic commissure, over the pons and lateral lobes of the cerebellum, was covered with a soft yellow exudate. The posterior surface of the spinal cord showed the same. The ventricles were much distended with a turbid fluid. There was suppuration of the mucous membrane of the nose, and of the ethmoidal and sphenoidal sinuses. Cultures showed the pneumococcus in the cerebro-spinal exudate, and from the nasal pus the pneumococcus and the staphylococcus aureus.

(c) *Primary pneumococcic meningitis*.—Future investigations will decide the proportion of cases of primary cerebro-spinal meningitis due to the pneumococcus and to the diplococcus intracellularis; and it will be an exceedingly interesting study to determine whether there are clinical differences such as separate so decidedly the meningitis complicating pneumonia from the cerebro-spinal fever. Judging from the bacteriological records given by Councilman, Mallory and Wright, and those of the Pathological Laboratory of the Johns Hopkins Hospital, the primary pneumococcic meningitis is rare in America—certainly much less infrequent than the secondary forms. Two of Councilman's ten cases, both infants, one of ten months, the other of six days, were primary. I cannot gather from Netter's statistics¹ how many of his cases were *primary*; nor without consulting the original can one say how many of the thirteen cases of sporadic meningitis studied by Malenchini (quoted by Netter) belonged to this group. Of our eight cases, as stated, only one was primary, and it is worth quoting:—

G. C., aged 44 (white), admitted at 6.30 p.m. on September 1, 1897, in an unconscious state. When seen by his brother on Saturday, the 29th, he was quite well. After going to bed on Saturday evening, the 30th, he had a severe chill. On Monday he was too ill to go to work. In the evening he became delirious, and ever since has had fever, has been restless, difficult to restrain, and very noisy. On admission the temperature was 105°; the pulse 136 and small. Dr. McCrae saw him a few minutes after he had been put to bed, and took very full and careful notes of the condition. The face was flushed and he looked distressed. There was no strabismus; the pupils were dilated and inactive. The head and neck were held stiffly, and the arms were held firmly flexed on the chest. There were no signs of pneumonia; the heart sounds were clear. The abdomen was not distended, there were no rose spots and the spleen could not be felt; there were no herpes. The skin had a diffuse mottling, but no purpuric spots were seen. There was no urethral discharge. The

¹ "Twentieth Century Practice," vol. xvi., p. 193.

leucocytes were 13,500 per cni., no alteration in a differential count of 500. The Widal reaction was not present. At 10 p.m. the temperature had reached 106° and he was given a bath at 80, with ice frictions, which reduced the rectal temperature to 103°. At 11 p.m. the body seemed to be held much more stiffly. At 12 midnight he was much worse, and at 12.40 a.m. he died, about three days and three hours from the time of the initial chill.

The *post mortem* (No. 989) showed a diffuse purulent meningitis of the hemispheres; the base much less involved and the ventricles not at all affected. The cord unfortunately was not examined. There was a recent diffuse nephritis, and the urine in the bladder contained albumen and tube cases. There was a general pneumococcus infection, no pneumonia or endocarditis, and the pneumococcus was separated in pure culture from the meningeal exudate.

V.—NOTE ON THE TREATMENT OF CEREBRO-SPINAL FEVER.

In our series of cases we have used no special drugs. Morphia has been freely given to control the pain. Ice sponging has been employed whenever the temperature reached above 102.5°. Our mortality, considering that the cases were as a rule quite severe, has been low, only eight of eighteen cases in the Hospital, and nine among the twenty-one cases which I have seen.

I have already spoken of the possible benefits in certain cases of the relief of pressure by the withdrawal of cerebro-spinal fluid.

In two of our cases the spinal canal has been opened, drained and irrigated. Winter (*Lancet*, 1891), in tuberculous meningitis, removed the spinal process of the second lumbar vertebra and drained the spinal canal. Von Ziemssen first attempted to use local therapeutics to the membranes of the cord by injecting a weak solution of iodine in a case of meningitis; and later Sahli practised permanent drainage through a canula and catheter. So far as I know, an extensive laminectomy had not been done for acute spinal meningitis until our first case, Nov. 6th, 1898, in which the operation was suggested and performed by Dr. Harvey W. Cushing, the first assistant in the surgical clinic of my colleague, Professor Halsted. The case has already been given in 'abstract under the section on Bacteriology' as the one in which the staphylococci only were present at the time of

¹ And reported in *Boston Medical and Surgical Journal*, December 29th, 1898.

operation. The spinal canal was thoroughly irrigated with salt solution and a quantity of purulent exudate washed out. No change followed in the existing paraplegia. The bladder and kidneys became infected, and he died about two months after the operation. The paraplegia persisted. At autopsy the spinal meninges were smooth and looked perfectly normal. It was impossible to say where the dura mater had been incised, and there were neither adhesions nor areas of the thickening on the pia-arachnoid. There were extensive changes in the cord itself and in nerve roots.

In case 12 laminectomy was performed on the fourth day by Dr. Cushing. The patient was very ill, and the spinal symptoms were especially pronounced. A catheter was passed beneath the dura mater, and the membranes drained and irrigated with salt solution. For several days he seemed much better. He developed a hæmorrhagic cystitis and pyelonephritis, and died on the sixth day after operation.

Dr. Musser, of Philadelphia, has also had an unsuccessful case. Dr. Rolleston and Mr. Herbert Allingham have reported¹ a case of sporadic cerebro-spinal meningitis, in which the patient recovered after laminectomy and drainage. On the principle of a desperate remedy for a desperate disease, the operation (which has been criticised adversely in an editorial note in the *Philadelphia Medical Journal*) seems justifiable in certain severe cases, in which, as in our first case, the spinal symptoms are very marked.

h
t.
er
s
y
l.
n
e
e